

3. Bhaskaran K, Hajat S, Haines A, Herrett E, Wilkinson P, Smeeth L. Effects of ambient temperature on the incidence of myocardial infarction. *Heart* 2009;95:1760–9.
4. Sue-Chu M. Winter sports athletes: long-term effects of cold air exposure. *Br J Sports Med* 2012;46:397–401.
5. Bougault V, Turmel J, St-Laurent J, Bertrand M, Boulet LP. Asthma, airway inflammation and epithelial damage in swimmers and cold-air athletes. *Eur Respir J* 2009;33:740–6.
6. Larsson K, Ohlén P, Larsson L, Malmberg P, Rydström PO, Ulriksen H. High prevalence of asthma in cross country skiers. *BMJ* 1993;307:1326–9.
7. Vergès S, Flore P, Blanchi MP, Wuyam B. A 10-year follow-up study of pulmonary function in symptomatic elite cross-country skiers—athletes and bronchial dysfunctions. *Scand J Med Sci Sports* 2004;14:381–7.
8. Nolan JP, Soar J, Zideman DA, et al., ERC Guidelines Writing Group. European Resuscitation Council Guidelines for Resuscitation 2010 Section 1. Executive summary. *Resuscitation* 2010;81:1219–76.
9. Parsons JP, Mastrorade JG. Exercise-induced bronchoconstriction in athletes. *Chest* 2005;128:3966–74.

Reply

We want to thank Drs. Chalkias and Xanthos from the University of Athens, Greece, for their interest in our study. Certainly, the mechanisms of cardiac arrest and a possible relationship with airway obstruction caused by exercise in cold weather are of interest in our northern part of the world, where many practice winter sports. There is a high prevalence of asthma among cross-country skiers.

We have statistics of temperatures from 10 long-distance (90 km) ski races with fatalities during the race and 114 races without fatalities. The mean morning temperature in races with fatal outcomes was -11.1°C , and average day temperature was -5.4°C (range $+0.8$ to -11.9°C). In the nonfatal races, the mean morning temperature was -9.6°C , and the average day temperature was -4.6°C (range $+5.3$ to -20.5°C). The variation between races

was great, and the lowest temperatures were recorded in races without fatal incidents.

From 1979 to 2012, 922 skiers were referred to a hospital emergency department. Of these, 20 skiers (2.2 %) had obstructive airway problems of asthma type. Thus, 1 of 43,000 cross-country skiers had to seek hospital care for airway obstruction.

In our original letter to the editor in the *Journal* (1), we described 20 cases of cardiac arrest. Interviews with the surviving patients and relatives of those who died did not reveal any history of obstructive airway symptoms.

In summary, our experience of cross-country skiing in Vasaloppet does not give any obvious support to airway obstruction as a major cause for morbidity during the race nor does it appear to be an important underlying mechanism for developing cardiac arrest during strenuous exercise in cold temperatures.

***Ulf Hållmarker, MD**
Karl Michaëlsson, MD, PhD
Johan Årnlöv, MD, PhD
Stefan James, MD, PhD

*Internal Medicine
 Mora lasarett Inst of Medicine
 Uppsala University
 794 92 Mora
 Sweden
 E-mail: ulf.hallmarker@ltdalarna.se

<http://dx.doi.org/10.1016/j.jacc.2012.10.022>

REFERENCE

1. Hållmarker U, Michaëlsson K, Årnlöv J, James S. Cardiac arrest in a long-distance ski race (Vasaloppet) in Sweden. *J Am Coll Cardiol* 2012;60:1431–2.